

POLYGONAL WORDS IN FREE GROUPS

SANG-HYUN KIM AND HENRY WILTON

[2000] Primary 20F36, 20F65 hyperbolic group, surface group

ABSTRACT. A longstanding question of Gromov asks whether every one-ended word-hyperbolic group contains a subgroup isomorphic to the fundamental group of a closed hyperbolic surface. An infinite family of word-hyperbolic groups can be obtained by taking doubles of free groups amalgamated along words that are not proper powers. We define the set of polygonal words in a free group of finite rank, and prove that polygonality of the amalgamating word guarantees that the associated square complex virtually contains a π_1 -injective closed surface. We provide many concrete examples of classes of polygonal words. For instance, in the case when the rank is two, we establish polygonality of words without an isolated generator, and also of almost all simple height-one words, including the Baumslag–Solitar relator $a^p(a^q)^b$ for $pq \neq 0$.

1. INTRODUCTION

A group G is called a *surface group* if G is isomorphic to the fundamental group of a closed surface S with non-positive Euler characteristic; if the Euler characteristic of S is negative, we say that G is a *hyperbolic surface group*. Motivated by the famous Surface Subgroup Conjecture in 3-manifold topology, Gromov asked the following question.¹

Question 1.1 (Gromov). Does every one-ended word-hyperbolic group contain a hyperbolic surface group?

In this paper, we introduce a new criterion to examine this question in the context of doubles of free groups. We denote by F the free group generated by $\{a_1, \dots, a_n\}$ for some $n > 1$. When the rank needs to be explicitly shown, we also denote F as F_n . We conventionally write the generators of F_2 as a and b . Let $w = v_1 \cdots v_l$ be a word in F where $v_i \in \{a_1^{\pm 1}, \dots, a_n^{\pm 1}\}$. Each v_i is called a *letter* of w . We say that w is *cyclically reduced* if $v_i \neq v_{i+1}^{-1}$ for each i , where the indices are taken modulo l . If w is cyclically reduced, we call l the *length* of w and write $|w| = l$. The word $w \in F$ is called *diskbusting* if $\{w\}$ does not belong to a proper free factor of F .

An infinite family of one-ended word-hyperbolic groups can be obtained by taking the *double* of F along w , namely $D(w) = F *_{\langle w \rangle} F$ where $w \in F$. The group $D(w)$ is one-ended if and only if w is diskbusting. Also, $D(w)$ is word-hyperbolic if and only if w is not a proper power [1]. The main result of [3] implies that

Kim supported by Basic Science Research Program (2010-0023515) and Mid-career Researcher Program (2010-0027001) through the National Research Foundation of Korea (NRF) funded by the MEST.

Wilton partially supported by NSF grant DMS-0906276.

¹Kahn and Markovic have recently announced a positive resolution of the Surface Subgroup Conjecture.

if $D(w)$ is word-hyperbolic and $H_2(G; \mathbb{Q}) \neq 0$, then $D(w)$ contains a hyperbolic surface group. Gordon and the second author proved a converse: the existence of a hyperbolic surface subgroup in $D(w)$ implies that $H_2(G; \mathbb{Q}) \neq 0$ for some finite-index subgroup G of $D(w)$ [4]. Using a certain Eilenberg–Mac Lane space $X(w)$ of $D(w)$ and looking at certain finite covers of $X(w)$, they also gave several sufficient conditions for $D(w)$ to have a finite-index subgroup G with $H_2(G; \mathbb{Q}) \neq 0$; see Section 2 for a precise definition of $X(w)$.

We will consider a combinatorial property, called *polygonality* of a word in F . A cyclically reduced word $w \in F$ is *polygonal* if there exists a surface S obtained by a side-pairing on polygonal disks $P_1, \dots, P_m$ where each edge in $S^{(1)}$ is oriented and labeled by $\{a_1, \dots, a_n\}$ in such a way that the image of each loop ∂P_i in S reads a non-trivial power of w and moreover, there do not exist two incoming edges or two outgoing edges of the same label at each vertex in $S^{(0)}$; see Definition 2.6. We say that a CW-complex X *virtually* contains a closed hyperbolic surface, if there is a homeomorphic embedding of a closed hyperbolic surface into a finite cover of X .

Theorem 2.11. *Let $w \in F$ be a cyclically reduced word.*

- (1) *If $w \in F$ is polygonal, then $X(w)$ virtually contains a closed hyperbolic surface.*
- (2) *If a closed hyperbolic surface embeds into a covering space of $X(w)$ with finitely generated fundamental group, then w is polygonal.*

Two words in F are *equivalent* if there exists an automorphism of F mapping one to the other. Theorem 2.11 will provide a way to detect surface subgroups in $D(w)$.

Corollary 2.14. *If $w \in F$ is equivalent to a polygonal word, then $D(w)$ contains a surface group.*

There are many examples of polygonal words. The following theorem summarizes the words in $F_2 = \langle a, b \rangle$ that we prove are polygonal. We use the notation $u^v = v^{-1}uv$ for words u, v . A word in F_2 of the form $w = \prod_{i=1}^l a^{p_i}(a^{q_i})^b$ is called a *simple height-one word* if all the integers p_i are of the same sign, all the integers q_i are of the same sign, and $l > 0$.

Theorem 1.2. *The following words in F_2 are polygonal.*

- (1) $w = \prod_i a^{p_i} b^{q_i}$ where $|p_i| > 1$ and $|q_i| > 1$ for each i (Theorem 4.6).
- (2) $w = \prod_i a^{p_i} b^{q_i}$ where $|p_i| > 1$ and $|q_i| = 1$ for each i , and $\sum_i (\text{sign}(p_i q_i) + \text{sign}(p_{i+1} q_i)) = 0$ (Theorem 4.7).
- (3) $w = \prod_i a^{p_i} (a^{q_i})^b$ where $|p_i| > 1$ and $|q_i| > 1$ for each i (Corollary 4.8).
- (4) A simple height-one word $w = \prod_i a^{p_i} (a^{q_i})^b$ where $pp' \leq q^2$ and $qq' \leq p^2$; here, we denote $p = \sum_i p_i, q = \sum_i q_i, p' = \sum_{|p_i|=1} 1$ and $q' = \sum_{|q_i|=1} 1$. (Theorem 5.6).

Theorem 1.2 (4) implies that *almost all* simple height-one words are polygonal, in a suitable sense (Remark 5.9).

A word $w \in F$ is *minimal* if no equivalent word to w is shorter. If one establishes polygonality (up to minimizing the length) for all diskbusting words then Question 1.1 would be answered affirmatively for doubles of free groups.

Conjecture 2.15 (Tiling Conjecture). *A minimal diskbusting word is polygonal.*

The hypothesis of minimality is necessary. There are examples of non-minimal diskbusting words that are not polygonal. In one sense, this is because surface subgroups need not manifest themselves as topological surfaces.

Proposition 3.2. *There exists a compact, non-positively curved square complex Y such that $\pi_1(Y)$ is a hyperbolic surface group but such that no subspace of Y is homeomorphic to a closed surface.*

Note. As this work was being written up, it came to our attention that N. Brady, M. Forester and E. Martinez-Pedroza have also, independently, developed a method of finding surface subgroups in certain graphs of free groups with cyclic edge groups. They use a theorem essentially the same as Theorem 4.6.

Acknowledgement. Both authors are deeply grateful for the numerous inspirations from Professor Cameron Gordon. Also, they are thankful for the guidance of Professor Alan Reid.

2. SURFACES IN A GRAPH OF GRAPHS WITH CYLINDRICAL EDGE SPACES

A *graph* means a 1-dimensional CW-complex. For a graph Γ , $V(\Gamma)$ and $E(\Gamma)$ will denote the vertex set and the edge set of Γ , respectively. If Γ_1 and Γ_2 are graphs, a continuous map $\phi: \Gamma_1 \rightarrow \Gamma_2$ sending each vertex to a vertex and each edge to an edge is called a *graph map*. An *immersion* is a graph map which is locally injective. $\text{Cay}(F)/F$ denotes the bouquet of n circles $\gamma_1, \dots, \gamma_n$ so that $\text{Cay}(F)/F$ is an Eilenberg–Mac Lane space for F . We always assume that each circle $\gamma_i: S^1 \rightarrow \text{Cay}(F)/F$ is equipped with an orientation and a label by a_i , and we identify $F = \pi_1(\text{Cay}(F)/F)$ by letting $[\gamma_i] = a_i$. We say that a loop $\gamma: S^1 \rightarrow \text{Cay}(F)/F$ reads $w = a_{i_1}^{m_1} a_{i_2}^{m_2} \cdots a_{i_l}^{m_l} \in F$ where $m_i \in \mathbb{Z}$, if γ is homotopic to the concatenation $\gamma_{i_1}^{m_1} \gamma_{i_2}^{m_2} \cdots \gamma_{i_l}^{m_l}$ as a based loop. For any cyclically reduced word $w \in F$, there exists an immersion $\gamma_w: S^1 \rightarrow \text{Cay}(F)/F$ reading w . Any immersion ϕ from a graph Γ to $\text{Cay}(F)/F$ induces a labeling and an orientation of each edge in Γ , so that ϕ preserves the labeling and the orientation. An edge in Γ labeled by a generator a_i is called an *a_i -edge*. For $F' \leq F$, $\text{Cay}(F)/F'$ will denote the coset Cayley graph of F/F' ; in other words, $\text{Cay}(F)/F'$ is the cover of $\text{Cay}(F, F)$ corresponding to F' .

We also consider an *abstract graph* in Scott–Wall’s sense [9]; that is, an abstract graph Γ consists of the vertex set $V(\Gamma)$, the edge set $E(\Gamma)$, the origin map $\iota: E(\Gamma) \rightarrow V(\Gamma)$, and a fixed-point-free involution on $E(\Gamma)$ sending e to $\bar{e}$. Note that $|E(\Gamma)|$ is twice the number of the edges in the geometric realization of Γ as a 1-dimensional CW-complex.

Definition 2.1. (1) Let Γ be an abstract graph, $\{G_v | v \in V(\Gamma)\}$ be a collection of finitely generated free groups and $\{w_e | e \in E(\Gamma)\}$ be a collection of non-trivial words such that $w_e \in G_{\iota(e)}$. We call the tuple $\mathcal{G} = (\Gamma, \{G_v | v \in V(\Gamma)\}, \{w_e | e \in E(\Gamma)\})$ a *graph of free groups with cyclic edge groups*.
 (2) Let $\mathcal{G} = (\Gamma, \{G_v | v \in V(\Gamma)\}, \{w_e | e \in E(\Gamma)\})$ be a graph of free groups with cyclic edge groups. Choose a graph X_v such that $\pi_1(X_v) = G_v$ and an immersion $\gamma_e: S^1 \rightarrow X_{\iota(e)}$ such that $[\gamma_e] = w_e \in G_{\iota(e)}$. For each pair $\{e, \bar{e}\}$, attach $S^1 \times [-1, 1]$ to $\coprod_v X_v$ such that $S^1 \times \{-1\}$ and $S^1 \times \{1\}$ are identified with γ_e and $\gamma_{\bar{e}}$, respectively; we denote by $X(\mathcal{G})$ the space thus obtained. Moreover, we let $\pi_1(\mathcal{G})$ denote the fundamental group of

$X(\mathcal{G})$. Note that $X(\mathcal{G})$ is well-defined up to homotopy equivalence, and so $\pi_1(\mathcal{G})$ is well-defined. We call each X_v a *vertex space* of $X(\mathcal{G})$, and the closure of each connected component of $X(\mathcal{G}) \setminus \coprod_v X_v$ a *cylinder* in $X(\mathcal{G})$. The image of each γ_e is called a *boundary component of the cylinder corresponding to $\{e, \bar{e}\}$* . Let $K \subseteq X(\mathcal{G}) \setminus \coprod_v X_v$ be the union of disjoint open annular neighborhoods of the cores of all the cylinders in $X(\mathcal{G})$. If C is a cylinder in $X(\mathcal{G})$ and H is a component of $C \setminus K$ intersecting with X_v , then we call H a *half-cylinder incident at X_v* .

Note that $X(\mathcal{G})$ is an Eilenberg–Mac Lane space for $\pi_1(\mathcal{G})$. The following lemma shows that any closed surface homeomorphically embedded in $X(\mathcal{G})$ is π_1 -injective.

Lemma 2.2. *Let $\mathcal{G}$ be a graph of free groups with cyclic edge groups and let S be a closed surface homeomorphically embedded into $X(\mathcal{G})$. Then:*

- (1) *S is a union of cylinders;*
- (2) *the inclusion $S \hookrightarrow X(\mathcal{G})$ induces a monomorphism $\pi_1(S) \hookrightarrow \pi_1(\mathcal{G})$;*
- (3) *S is of non-positive Euler characteristic.*

Proof. (1) Suppose there exists $s \in S$ such that for any cylinder C containing s , S does not intersect $\text{int}(C)$. Let $D \approx D^2$ be a Euclidean neighborhood of s in S which is so small that D intersects only the cylinders that contains s . Then $D \subseteq S$ does not intersect the interiors of any cylinders, by the assumption on s . It follows that $D \subseteq X(\mathcal{G})^{(1)}$, which is a contradiction.

Hence, for any $s \in S$ there exists a cylinder C such that $s \in C$ and $S \cap \text{int}(C) \neq \emptyset$. In other words, $S \subseteq \cup_{S \cap \text{int}(C) \neq \emptyset} C$. Fix any cylinder C such that $S \cap \text{int}(C) \neq \emptyset$. Since $\text{int}(C)$ is an open cylinder, any sufficiently small regular neighborhood $V \subseteq X(\mathcal{G})$ of $t \in S \cap \text{int}(C)$ is actually Euclidean and so, $V \subseteq S$. This means $S \cap \text{int}(C)$ is open in $\text{int}(C)$. On the other hand, compactness of S implies $S \cap \text{int}(C)$ is a closed subset of $\text{int}(C)$. So, $S \cap \text{int}(C) = \text{int}(C)$. From $\cup_{S \cap \text{int}(C) \neq \emptyset} \text{int}(C) \subseteq S \subseteq \cup_{S \cap \text{int}(C) \neq \emptyset} C$, we have $S = \cup_{S \cap \text{int}(C) \neq \emptyset} C$.

(2) Let X_0 be the union of all the vertex spaces of $X(\mathcal{G})$ and let $S_0 = S \cap X_0$. Let Γ be the underlying graph of $\mathcal{G}$. Consider S as a graph of spaces where each connected component of S_0 is a vertex space, and each cylinder is an edge space, such that the underlying graph is a subgraph of Γ . This gives $\pi_1(S)$ the structure of a graph of groups $\mathcal{K}$. Any element of $\pi_1(S)$ that is in reduced form as an element of $\pi_1(\mathcal{G})$ is also in reduced form with respect to $\mathcal{K}$. Hence the natural map $\pi_1(S) \rightarrow \pi_1(\mathcal{G})$ is injective.

(3) This is immediate from part (1). $\square$

Remark 2.3. In the proof of Lemma 2.2 (2), we have only used the assumption that S is a connected union of cylinders, rather than a closed surface.

As mentioned in the introduction, we are most concerned with the *double* of a free group along a word: this is the a special case of a graph of free groups with cyclic edge groups $\mathcal{G} = (\Gamma, \{G_v\}, \{w_e\})$ in which the underlying graph has two vertices and a single edge $\{e, \bar{e}\}$, and there is an isomorphism from $G_{\iota(e)}$ to $G_{\iota(\bar{e})}$ mapping w_e to $w_{\bar{e}}$. We write $X(w) = X(\mathcal{G})$.

Let $\gamma_w: S^1 \rightarrow \text{Cay}(F)/F$ be an immersed loop reading w . By convention, we will always choose $X(w)$ to be the 2-dimensional square complex obtained by taking two copies of $\text{Cay}(F)/F$ and gluing a cylinder between the two copies of $\gamma_w \subseteq \text{Cay}(F)/F$. Given a word $w \in F$ and a finite-index subgroup $F' \leq F$, there

is a particular finite cover $Y(w, F')$ of $X(w)$, considered in [4]. That is, consider the *pullback* $\tilde{\gamma}_w: T \rightarrow S^1$ of γ_w along the covering map $\text{Cay}(F)/F' \rightarrow \text{Cay}(F)/F$, yielding the commutative diagram below. The space T is a finite-sheeted covering space of S^1 , and hence a disjoint union of circles. Then, $Y(w, F')$ is defined to be the graph of spaces corresponding to the first row of the diagram. Note that $X(w)$ is the graph of spaces corresponding to the second row. There is a natural covering map $Y(w, F') \rightarrow X(w)$ of degree $[F : F']$.

$$\begin{array}{ccccc} \text{Cay}(F)/F' & \xleftarrow{\tilde{\gamma}_w} & T = \coprod S^1 & \xrightarrow{\tilde{\gamma}_w} & \text{Cay}(F)/F' \\ \downarrow & & \downarrow & & \downarrow \\ \text{Cay}(F)/F & \xleftarrow{\gamma_w} & S^1 & \xrightarrow{\gamma_w} & \text{Cay}(F)/F \end{array}$$

A restriction of $\tilde{\gamma}_w$ to a connected component of T is called an *elevation* of γ_w . Elevations can be interpreted algebraically:

Lemma 2.4 ([11], Lemma 2.7). *There is a natural bijection between the set of elevations of γ_w and the set of double cosets*

$$F' \backslash F / \langle w \rangle.$$

For each double coset $F'g\langle w \rangle$, the corresponding elevation $\tilde{\gamma}_w$ is freely homotopic to the lifting of $\gamma_g \gamma_w^{n_g} \gamma_g^{-1}$ with respect to $\text{Cay}(F)/F' \rightarrow \text{Cay}(F)/F$, where n_g is the minimal positive integer such that $gw^{n_g}g^{-1} \in F'$.

Lemma 2.5. *Let w be a cyclically reduced word in F . If $X(w)$ virtually contains a closed hyperbolic surface, then $Y(w, F')$ contains a homeomorphically embedded closed hyperbolic surface for some $[F : F'] < \infty$.*

Proof. Suppose $X(\mathcal{G})$ is a cover of $X(w)$ containing a homeomorphically embedded closed hyperbolic surface S where $\mathcal{G}$ is a graph of free groups with cyclic edge groups. By Lemma 2.2, S is a union of cylinders in $X(\mathcal{G})$. Let K denote the union of disjoint open annular neighborhoods of the cores of the cylinders in $X(\mathcal{G})$ and let S_0 be a connected component of $S \setminus K$ such that $\chi(S_0) < 0$. Then S_0 is contained in the neighborhood N of some vertex space X' of $X(\mathcal{G})$. Since $X' \rightarrow \text{Cay}(F)/F$ is a finite cover, $X' = \text{Cay}(F)/F'$ for some $[F : F'] < \infty$. Let $\gamma_w: S^1 \rightarrow \text{Cay}(F)/F$ denote the immersed loop reading w . Recall that $Y(w, F')$ is obtained by taking two copies of $X' = \text{Cay}(F)/F'$ and gluing cylinders along the copies of the connected components of the pullback $\tilde{\gamma}_w$ of γ_w along $\text{Cay}(F)/F' \rightarrow \text{Cay}(F)$. As S_0 is the union of some half-cylinders incident at X' , which are glued along some elevations of γ_w , the double S of S_0 glued along ∂S_0 is a closed hyperbolic surface in $Y(w, F')$. $\square$

A *polygonal disk* P is a 2-dimensional disk such that ∂P comes with the CW-structure of a polygon. If the edges (vertices, respectively) of P are denoted as $x_1, x_2, \dots, x_m$, we will consider the indices as taken to be modulo m , and will implicitly assume that x_i and x_{i+1} intersect with a common vertex (edge, respectively). We let $E(\partial P)$ denote the set of 1-cells (edges) in ∂P . A *pairing* on a set X is an equivalence relation such that each equivalence class consists of exactly two elements. A *partial side-pairing* on a collection of polygonal disks $P_1, \dots, P_m$ is a pairing $\sim$ on a subset of $\coprod_i E(\partial P_i)$ along with a choice of a homeomorphism identifying the edges related by $\sim$. A *side-pairing* refers to a partial side-pairing which

is defined on the whole set $\coprod_i E(\partial P_i)$. For a (partial, respectively) side-pairing $\sim$ on $P_1, \dots, P_m$, we denote by $\coprod_i P_i / \sim$ the closed (bounded, respectively) surface obtained from $\coprod_i P_i$ by identifying edges in $\coprod_i \partial P_i$ through $\sim$. For $S = \coprod_i P_i$, note that there is a natural cellular map $P_i \rightarrow S$ so that S has the structure of a 2-dimensional CW-complex. We will let $m(S)$ denote the number of 2-cells in S .

Definition 2.6. Let w be a cyclically reduced word in F .

- (1) Suppose there exist a partial side-pairing $\sim$ on some polygonal disks $P_1, \dots, P_m$, and an immersion $\phi: S^{(1)} \rightarrow \text{Cay}(F)/F$ where $S = \coprod_i P_i / \sim$ such that the composition $\partial P_i \rightarrow S^{(1)} \rightarrow \text{Cay}(F)/F$ is an immersion reading a non-zero power of w for each i . Then we say that S is a w -polygonal surface (with respect to ϕ).
- (2) We say that w is *polygonal*, if there exists a closed w -polygonal surface S such that $\chi(S) < m(S)$.

The most convenient way to specify ϕ is to label and orient the edges of the boundaries ∂P_i according to their images under ϕ . Let P be a polygonal disk equipped with an orientation and a label by $\{a_1, \dots, a_n\}$ for each edge of ∂P . We say that ∂P reads w (or, simply P reads w) if the orientation and the label preserving graph map $\partial P \rightarrow \text{Cay}(F)/F$ reads w . If P is a polygonal disk whose boundary reads w and u is a subword of w , the interval in ∂P reading u will be simply referred to as u when there is no danger of confusion. We will also view a w -polygonal surface S as obtained by a side-pairing $\sim$ on polygonal disks $P_1, \dots, P_m$ such that each ∂P_i reads a power of w and $\sim$ respects orientations and labels of the edges; moreover, one requires that at each vertex of S there are no two incoming edges or two outgoing edges of the same label, and $\chi(S) < m(S)$.

Remark 2.7. Let $w \in F$ be a cyclically reduced word which is not a proper power, and S be a closed connected w -polygonal surface. Note that the condition $\chi(S) < m(S)$ is equivalent to forbidding the following three cases:

- (i) $m(S) = 1$ and $S \approx S^2$,
- (ii) $m(S) = 2$ and $S \approx S^2$, or
- (iii) $m(S) = 1$ and $S \approx \mathbb{R}P^2$.

The case (i) occurs when $S = (P/\sim) \approx S^2$ for some side-pairing $\sim$ on a polygonal disk P . Then there must exist two consecutive edges of ∂P which are identified by $\sim$ such that $\sim$ fixes a common vertex. This is impossible by the definition of an immersion. The case (ii) is equivalent to the condition that S consists of two identical polygonal disks whose boundaries (via immersions into $\text{Cay}(F)/F$) both read the same power w^i , and $\sim$ identifies each pair of edges corresponding to the same letter of w^i . The case (iii) amounts to saying that S consists of a polygonal disk P such that an immersion $\phi: \partial P \rightarrow \text{Cay}(F)/F$ reads $w^k = u^2$ for some $k > 0$ and $u \in F$, and $\sim$ identifies each pair of edges on ∂P by a π -rotation; this means, the i -th letter of u^2 is identified with the $(i + |u|)$ -th letter of u^2 . For example, take $k = 2$ and $u = w$. As (i), (ii) and (iii) occur only in these obvious cases, we will often omit the proof that $\chi(S) < m(S)$ for a w -polygonal surface S when establishing polygonality of w .

Example 2.8. If $w \in F$ is a surface group relator such as $\prod_i [a_{2i-1}, a_{2i}]$ or $\prod_i a_i^2$, then w is polygonal. This is because one can take a polygonal disk whose boundary reads w and glue the edges of the same labels respecting the orientations.

Lemma 2.9. *Suppose $w \in F$ is cyclically reduced and $S = \coprod_i P_i / \sim$ is a closed connected w -polygonal surface such that $\chi(S) < m(S)$. Suppose Q and Q' are distinct vertices of ∂P_i for some i and one of the two intervals on ∂P_i cut by $\{Q, Q'\}$ reads a power of w . Then Q is not identified to Q' by $\sim$.*

Proof. Let $\phi: S^{(1)} \rightarrow \text{Cay}(F)/F$ be the immersion with respect to which S is w -polygonal. Name the vertices of P_i as $v_1, \dots, v_{kl}$ where $k > 0$ and $l = |w|$. We may assume $Q = v_1$ and $Q' = v_{hl+1}$ for some $0 < h < k$. Suppose $v_1 \sim v_{hl+1}$. Since the orientations and the labels of the edges $(v_1, v_2), (v_2, v_3), \dots$ coincide with those of $(v_{hl+1}, v_{hl+2}), (v_{hl+2}, v_{hl+3}), \dots$ respectively, the immersion condition of $\partial P_i \rightarrow S^{(1)} \rightarrow \text{Cay}(F)/F$ implies that $v_2 \sim v_{hl+2}, v_3 \sim v_{hl+3}, \dots$. This is possible only when $\sim$ pairs all the edges of ∂P_i by a π -rotation; hence, $m(S) = 1 = \chi(S)$. $\square$

A graph Γ embedded in a compact surface with boundary S' is a *spine* if the closure of every connected component of $S' \setminus \Gamma$ is an annulus, with one boundary component contained in Γ and one boundary component contained in $\partial S'$.

Lemma 2.10. *Let w be a cyclically reduced word in F . Then w is polygonal if and only if there exists a commutative diagram*

$$\begin{array}{ccc}
 T = \coprod S^1 & \xrightarrow{\tilde{\gamma}_w} & \text{Cay}(F)/F' \\
 \downarrow & \swarrow \text{incl.} & \nearrow \tilde{\psi} \\
 & \partial S' \hookrightarrow S' & \\
 \downarrow & \swarrow & \searrow \\
 S^1 & \xrightarrow{\gamma_w} & \text{Cay}(F)/F
 \end{array}$$

such that $[F : F'] < \infty$, γ_w is an immersion reading w , $\tilde{\gamma}_w$ is the pullback of γ_w along $\text{Cay}(F)/F' \rightarrow \text{Cay}(F)/F$, S' is a hyperbolic surface with boundary, $\tilde{\psi}$ is an embedding when restricted to a spine of S' and $\partial S' \rightarrow T = \coprod S^1$ is an inclusion.

Proof. ($\Rightarrow$) Choose a closed connected w -polygonal surface $S = \coprod P_i / \sim$ with respect to an immersion $\phi: S^{(1)} \rightarrow \text{Cay}(F)/F$ such that $\chi(S) < m(S)$. Since F is subgroup separable, the immersion ϕ lifts to an embedding $\tilde{\phi}: S^{(1)} \rightarrow \text{Cay}(F)/F'$ for some $[F : F'] < \infty$ [10]. Let S' be obtained by drilling a hole in the interior of each of P_i 's. Then $\chi(S') = \chi(S) - m(S) < 0$ and $S^{(1)}$ is a spine of S' . Define $\tilde{\psi}$ to be the composition $S' \xrightarrow{\sim} S^{(1)} \rightarrow \text{Cay}(F)/F'$ so that $\tilde{\psi}_*: \pi_1(S') \rightarrow F'$ is injective. Because the boundary of each P_i reads a power of w , there is a covering map $\partial S' \rightarrow S^1$ whose composition with γ_w equals the composition $\partial S' \hookrightarrow S' \rightarrow \text{Cay}(F)/F' \rightarrow \text{Cay}(F)/F$.

Let $T = \coprod S^1$ be the pullback of γ_w along the covering map $\text{Cay}(F)/F' \rightarrow \text{Cay}(F)/F$. By the universal property, the covering map $\partial S' \rightarrow S^1$ lifts to $q: \partial S' \rightarrow T$. Let $\partial_i S'$ denote the boundary component of S' drilled on P_i . The restriction of q to $\partial_i S'$ is a finite covering map onto its image, say $A \subseteq T$. $A \rightarrow S^1 \xrightarrow{\gamma_w} \text{Cay}(F)/F$ reads w^r for some $r > 0$. If $\partial_i S' \rightarrow A$ is a k -to-1 map for some $k > 1$, then $\partial P_i \rightarrow \text{Cay}(F)/F$ will read $(w^r)^k$. In particular, $\partial P_i \rightarrow S^{(1)} \rightarrow \text{Cay}(F)/F'$ maps a proper subinterval on ∂P_i onto the image of $A \rightarrow \text{Cay}(F)/F'$. This is a contradiction to Lemma 2.9, for $A \rightarrow \text{Cay}(F)/F' \rightarrow \text{Cay}(F)/F$ reads w^r . Hence the restriction of q to each boundary component of S' is a homeomorphism onto

the image. Suppose $\partial_i S'$ and $\partial_j S'$ map onto the same component of T for $i \neq j$. Then $\partial P_i \rightarrow S^{(1)} \hookrightarrow \text{Cay}(F)/F'$ and $\partial P_j \rightarrow S^{(1)} \hookrightarrow \text{Cay}(F)/F'$ have the same image. It follows that $P_i \cup P_j$ maps to a 2-sphere in S . Since we have excluded the case $\chi(S) = m(S) = 2$, we conclude that $q: \partial S' \rightarrow T$ must be an embedding.

($\Leftarrow$) Take Γ to be the spine for S' provided by the hypotheses. For each component $\partial_i S'$ of $\partial S'$, let A_i be the closure of the component of $S' \setminus \Gamma$ that contains $\partial_i S'$. Glue disks Q_i to $\partial_i S'$, to form a closed surface S . Let $P_i = Q_i \cup A_i$, a disk with boundary contained in Γ ; give S the CW-structure in which $S^{(1)} = \Gamma$ and the P_i are the 2-cells. Because $\partial Q_i = \partial_i S'$ reads a power of w and A_i is an annulus, it follows that each ∂P_i also reads a power of w under the composition $\Gamma \rightarrow \text{Cay}(F)/F' \rightarrow \text{Cay}(F)/F$. Note that $\chi(S) - m(S) = \chi(S') < 0$, where $m(S)$ is the number of 2-cells P_i . $\square$

Theorem 2.11. *Let $w \in F$ be a cyclically reduced word.*

- (1) *If $w \in F$ is polygonal, then $X(w)$ virtually contains a closed hyperbolic surface.*
- (2) *If a closed hyperbolic surface embeds into a covering space of $X(w)$ with the finitely generated fundamental group, then w is polygonal.*

Proof. Let $\gamma_w: S^1 \rightarrow \text{Cay}(F)/F$ be an immersion reading w .

(1) There exist a hyperbolic surface S' and a finite-index subgroup $F' \leq F$ satisfying the commutative diagram in Lemma 2.10, where $\tilde{\gamma}_w: T = \coprod S^1 \rightarrow \text{Cay}(F)/F'$ denotes the pullback of γ_w along $\text{Cay}(F)/F' \rightarrow \text{Cay}(F)/F$. Let S_0 be the closed surface obtained by taking the double of S' along $\partial S'$. The composition $\partial S' \hookrightarrow T = \coprod S^1 \rightarrow \text{Cay}(F)/F'$ restricts to distinct elevations on the components of $\partial S'$. By the definition of $Y(w, F')$, S_0 is homeomorphic to a union of distinct cylinders in $Y(w, F')$. Note $\chi(S_0) = 2\chi(S') < 0$.

(2) Since $D(w)$ is the fundamental group of a graph of free groups with cyclic edge groups which is word-hyperbolic, $D(w)$ is subgroup separable [12]. If a closed hyperbolic surface S_0 homeomorphically embeds into a covering space of $X(w)$ with finitely generated fundamental group, then S_0 homeomorphically embeds also into a finite cover of $X(w)$ [8]. By Lemma 2.5, S_0 may be assumed to homeomorphically embed into $Y(w, F')$ for some $[F : F'] < \infty$. By Lemma 2.2, S_0 is a union of cylinders. Let m denote the number of cylinders in S . Pick one of the two vertex spaces in $Y(w, F')$ and name it simply as $\text{Cay}(F)/F'$. Let K be the disjoint union of the open annular neighborhoods of the cores of the cylinders in $Y(w, F')$. Define Y' (S' , respectively) to be the connected component of $Y \setminus K$ ($S_0 \setminus K$, respectively) intersecting with $\text{Cay}(F)/F'$. Note that $\Gamma = S' \cap \text{Cay}(F)/F'$ is a spine for S' . Let $\tilde{\gamma}_w: T = \coprod S^1 \rightarrow \text{Cay}(F)/F'$ denote the pullback of γ_w along $\text{Cay}(F)/F' \rightarrow \text{Cay}(F)/F$. Then Y' is the union of $\text{Cay}(F)/F'$ and the half-cylinders glued along the restrictions of $\tilde{\gamma}_w$ to the components of T . Moreover, S' is some collection of those half-cylinders and so, there exist an inclusion $\partial S' \hookrightarrow T$ yielding the commutative diagram in Lemma 2.10. S' is hyperbolic since $\chi(S') = \frac{1}{2}\chi(S_0) < 0$. $\square$

Remark 2.12. Let S be a closed connected w -polygonal surface. From the proof of Theorem 2.11, one can actually find the degree of a cover $Y(w, F') \rightarrow X(w)$ where $Y(w, F')$ contains an embedded closed hyperbolic surface S_0 . Namely, since the immersion $S^{(1)} \rightarrow \text{Cay}(F)/F$ lifts to an embedding $S^{(1)} \rightarrow \text{Cay}(F)/F'$ where

the embedding does not increase the number of vertices [10],

$$\deg(Y(w, F') \rightarrow X(w)) = \deg(\text{Cay}(F)/F' \rightarrow \text{Cay}(F)/F) = |(\text{Cay}(F)/F')^{(0)}| = |S^{(0)}|.$$

Moreover, one can specify the Euler characteristic of S_0 as $\chi(S_0) = 2(\chi(S) - m(S))$.

Example 2.13. In an earlier version of [4], it was asked whether $D(w)$ contains a surface group for $w = a^{-2}b^{-1}a^{-1}bab^{-1}ab \in F_2$. Figure 1 describes a side-pairing $\sim$ of a polygonal disk P with its boundary reading w^2 so that $S = P/\sim$ is a closed surface of Euler characteristic -1 . Note that no vertex of S has two incoming or two outgoing edges of the same label. By Remark 2.12, one can find a closed surface S' in $Y(w, F')$ for some $[F : F'] = |S^{(0)}| = 7$ such that $\chi(S') = 2(\chi(S) - m(S)) = 2(-1 - 1) = -4$. In particular, $D(w)$ contains a surface group. One can similarly show that $w = a^p(a^{-1})^b a^q a^b$ is polygonal for any $pq \neq 0$.




FIGURE 1. A side-pairing on a polygonal disk P . Single and double arrows denote a -edges and b -edges, respectively. In (a), starting from the vertex marked by A , following the dashed arrow will read $w^2 = (a^2(a^{-1})^b aa^b)^2$. Let $\sim$ glue a pair of intervals, marked as $\{7, 8, 9\}$ and $\{7', 8', 9'\}$, so that P becomes an annulus as shown in (b). In (b), let a side-pairing $\sim$ further pair the edges marked by $1, \dots, 6$ with the edges marked by $1', \dots, 6'$, respectively. Now $S = P/\sim$ is a closed w -polygonal surface of Euler characteristic -1 .

Corollary 2.14. *If $w \in F$ is equivalent to a polygonal word, then $D(w)$ contains a surface group.*

Proof. We may assume w is polygonal. By Theorem 2.11, $X(w)$ virtually contains a closed surface S . Lemma 2.2 implies that $\pi_1(S)$ embeds into $D(w) = \pi_1(X(w))$. $\square$

We note that a polygonal word is not an element of a basis; for otherwise, $D(w)$ would be free.

Conjecture 2.15 (Tiling Conjecture). *A minimal diskbusting word is polygonal.*

3. NON-POLYGONAL WORDS

The following example shows that the hypothesis of minimality is necessary in the Tiling Conjecture.

Example 3.1. There are diskbusting words which are not polygonal. For example, one can show that $w = abab^2ab^3$ is not polygonal as follows. Suppose there exists a w -polygonal surface S such that $\chi(S) < m(S)$. Then there exists a vertex v in $S^{(0)}$ such that $\deg(v)$ is at least 3. From the fact that w is positive, it follows that $\deg(v)$ is even. So $\deg(v) = 4$, and in particular, there exists an incoming a -edge at v . Since the letter a is always followed by b in w and exactly one b -edge is outgoing at v , $\deg(v) = 2$; hence, we have a contradiction. Note that by the automorphism $a \mapsto ab^{-2}$ and $b \mapsto b$, w maps to $a(a^2)^b$. In Example 5.7, $a(a^2)^b$ is shown to be polygonal. It is not hard to see that $a(a^2)^b$ is minimal.

So we see that a double may contain a surface group which does not manifest itself as an embedded surface in a cover. One can interpret this in terms of the existence of compact complexes which are homotopic to a surface but which do not contain embedded surfaces, as the next proposition shows. We refer the reader to [2] for the definition and properties of CAT(0) spaces. A square complex is non-positively curved (meaning that its universal cover is CAT(0)) if and only if the link of every vertex contains no cycles of length less than four, by Gromov's Link Condition. In particular, the complex $X(w)$ is a non-positively curved square complex.

Proposition 3.2. *There exists a compact, non-positively curved square complex Y such that $\pi_1(Y)$ is a hyperbolic surface group but such that no subspace of Y is homeomorphic to a closed surface.*

Proof. Let w be a diskbusting, non-polygonal word that is equivalent to a polygonal one, as in Example 3.1. Then $D(w)$ contains a surface subgroup H . This subgroup is quasiconvex (this follows, for instance, from the fact that H is a virtual retract [11]). The universal cover $\tilde{X}$ of $X(w)$ is a Gromov-hyperbolic CAT(0) square complex, quasi-isometric to the Cayley graph of $D(w)$ by the Švarc–Milnor Lemma. Because every quasigeodesic in a Gromov-hyperbolic metric space is uniformly close to a geodesic, every orbit of H is quasiconvex in the 1-skeleton of $\tilde{X}$. Therefore, by [5, Theorem H], it follows that H acts cocompactly on a convex subcomplex $\tilde{Y}$ of $\tilde{X}$.

Let $Y = \tilde{Y}/H$. This is a compact, non-positively curved square complex with fundamental group H . However, the inclusion $\tilde{Y} \hookrightarrow \tilde{X}$ descends to an embedding $Y \hookrightarrow \tilde{X}/H$, a covering space of $X(w)$ with fundamental group H . Therefore, by Theorem 2.11, no compact subspace of Y is homeomorphic to a compact surface. $\square$

Remark 3.3. As this article was in its final stages of revision, we heard of a simpler construction of a similar example, due to Noel Brady [6].

4. WORDS WITHOUT ISOLATED GENERATORS

We consider words of the following form in $F_n = \langle a_1, \dots, a_n \rangle$:

$$w = \prod_{i=1}^l a_{k_i}^{p_i}, \quad \text{where } l > 0, k_i \neq k_{i+1} \text{ and } |p_i| > 1.$$

Note that we take the index i modulo l . Such w is said to have *no isolated generators*.

Throughout this section, we will employ the following notations. Take two identical polygonal disks P and P' such that ∂P and $\partial P'$ read w . Put $p = \sum_{i=1}^l |p_i|$. Label the edges of the p -gon ∂P as $e_1, e_2, \dots, e_p$. Fix an orientation-preserving homeomorphism $h: P \rightarrow P'$ such that e_j and $h(e_j)$ have the same label and the orientation. Consider a partial side-pairing $\sim$ on $P \amalg P'$ such that $e_j \sim h(e_{j+1})$ if and only if the label and the orientation of e_j coincide with those of $h(e_{j+1})$; see Figure 2 (a). For each $1 \leq i \leq l$, there exists a boundary component ∂_i of $S = P \amalg P' / \sim$ such that ∂_i is a 2-cycle whose edges are labeled by a_{k_i} and $a_{k_{i+1}}$ (Figure 2 (b)).

Let $\rho_i = (\text{sign}(p_i)k_i, \text{sign}(p_{i+1})k_{i+1})$ for each $1 \leq i \leq l$. Recall from Section 2 that an edge of $S^{(1)}$ which is labeled by a_k is called an a_k -edge. If $\text{sign}(p_i) = 1$ then there exists an incoming a_{k_i} -edge at ∂_i , and if $\text{sign}(p_{i+1}) = -1$ then there exists an incoming $a_{k_{i+1}}$ -edge at ∂_i . Likewise, if $\text{sign}(p_i) = -1$ then there exists an outgoing a_{k_i} -edge at ∂_i , and if $\text{sign}(p_{i+1}) = 1$ then there exists an outgoing $a_{k_{i+1}}$ -edge at ∂_i .

Let $V_n = \{\pm 1, \pm 2, \dots, \pm n\}$ and let

$$E_n = \{(i, j) \mid i, j \in V_n \text{ and } |i| \neq |j|\} \subseteq V_n \times V_n.$$

Define $\mathcal{A}_n^*$ to be the free abelian monoid on the set E_n , and let $\mathcal{A}_n$ be the quotient of $\mathcal{A}_n^*$ by the relation $(i, j) = (-j, -i)$. Then $\mathcal{A}_n$ is also a free abelian monoid. Put $\rho(w) = \sum_{i=1}^l \rho_i \in \mathcal{A}_n$. Define $\mathcal{T}_n \leq \mathcal{A}_n$ by

$$\mathcal{T}_n = \langle \{(c_1, c_2) + (c_2, c_3) + \dots + (c_r, c_1) \mid c_1, c_2, \dots, c_r \in V_n \text{ are all distinct}\} \rangle.$$




FIGURE 2. An example of S when $w = a_1^3 a_2^2 a_1^{-2} a_2^{-3}$. Single and double arrows denote a_1 -edges and a_2 -edges, respectively. (a) Rotate $\partial P'$ by $\frac{2\pi}{10}$ and identify the overlapping edges if the labels and the orientations coincide. (b) Edges intersecting with a boundary component ∂_j of $S = P \amalg P' / \sim$. In this figure, $\rho_j = (-1, 2)$.

Theorem 4.1. *Let $w = \prod_{i=1}^l a_{k_i}^{p_i} \in F_n$ where $l > 0, k_i \neq k_{i+1}$ and $|p_i| > 1$. If $\rho(w) \in \mathcal{T}_n$, then w is polygonal.*

Proof. We may assume $l > 1$, as proper powers are polygonal by definition. We will define a side-pairing $\sim'$ on $P \amalg P'$ such that $\sim'$ is an extension of $\sim$. Put $p = \sum_{i=1}^l |p_i|$. Since $\rho(w) = \sum_{i=1}^l \rho_i \in \mathcal{T}_n$, we can write $\{1, 2, \dots, p\} = \coprod_{h=1}^q A_h$ so that for each h , $\sum_{i \in A_h} \rho_i$ is a generator of $\mathcal{T}_n$. Choose any $1 \leq h \leq q$ and write $A_h = \{i_1, \dots, i_r\}$. We will define $\sim'$ for the edges of $\partial_{i_1} \cup \dots \cup \partial_{i_r}$. We have $\rho_{i_1} + \rho_{i_2} + \dots + \rho_{i_r} = (c_1, c_2) + (c_2, c_3) + \dots + (c_r, c_1)$ for some distinct $c_1, c_2, \dots, c_r \in V_n$. Note that at a vertex of ∂_{i_j} , an $a_{|c_j|}$ -edge is incoming or outgoing according to whether $\text{sign}(c_j) = 1$ or $\text{sign}(c_j) = -1$. Also, an $a_{|c_{j+1}|}$ -edge intersects ∂_{i_j} in a similar fashion; see Figure 3. Let $\sim'$ glue the $a_{|c_j|}$ -edge of $\partial_{i_{j-1}}$ to the $a_{|c_j|}$ -edge of ∂_{i_j} , for $j = 1, 2, \dots, r$. At any vertex of $\partial_{i_1} \cup \dots \cup \partial_{i_r} / \sim'$, there do not exist two incoming or two outgoing edges of the same label, since $|c_j|$'s are all distinct. As $1 \leq h \leq q$ was arbitrarily chosen, $\sim'$ is defined for all the edges of $\partial S = \partial(P \amalg P' / \sim)$. We can consider $\sim'$ as an extension of $\sim$ such that $P \amalg P' / \sim'$ is a closed w -polygonal surface. $\square$




FIGURE 3. $\rho_{i_1} + \rho_{i_2} + \rho_{i_3}$ is a generator of $\mathcal{T}_n$. The single, double and the triple arrows denote edges labeled by a_1, a_2 and $a_3 \in F_n$, respectively. Note that there exists exactly one incoming a_i -edge and one outgoing a_i -edge at each vertex of $\partial_{i_1} \cup \partial_{i_2} \cup \partial_{i_3}$, for each $i = 1, 2, 3$.

Remark 4.2. (1) In Theorem 4.1, $\mathcal{A}_n$ and $\mathcal{T}_n$ are independent of the choices of w . Also, $\rho(w)$ can be easily computed as

$$\rho(w) = \sum_i (\text{sign}(p_i) a_{k_i}, \text{sign}(p_{i+1}) a_{k_{i+1}}).$$

(2) In Theorem 4.1, the hypothesis that $\rho(w) \in \mathcal{T}_n$ is necessary. For example, $w = a^2 b^2 c^3 b^{-3} \in \langle a, b, c \rangle = F_3$ does not have any isolated generators. But

$$\langle w, a, b^2 c b^{-2} \rangle = \langle b^2 c^3 b^{-3}, a, b^2 c b^{-2} \rangle = \langle b^{-1}, a, b^2 c b^{-2} \rangle = \langle a, b, c \rangle,$$

It follows that w is an element of a basis, and hence not polygonal.

Corollary 4.3. Let $w = \prod_{i=1}^l a_{k_i}^{p_i} \in F_n$ for $l > 0, k_i \neq k_{i+1}$ and $|p_i| > 1$. If k_i 's are all distinct, then w is polygonal.

Proof. Since a_{k_i} 's are all distinct, $\rho(w) = \sum_i (\text{sign}(p_i) k_i, \text{sign}(p_{i+1}) k_{i+1})$ is a generator of $\mathcal{T}_n$. $\square$

Example 4.4. (1) $w = a_1^2 a_2^{-3} a_3^2 \in F_3$ is polygonal, by Corollary 4.3.

- (2) Let $w = a_1^6 a_2^{-3} a_3^5 a_2^4 a_3^{-7} \in F_3$. We have $\rho(w) = (1, -2) + (-2, 3) + (3, 2) + (2, -3) + (-3, 1) = (1, -2) + (-2, 3) + (-2, -3) + (3, -2) + (-3, 1) = ((1, -2) + (-2, -3) + (-3, 1)) + ((-2, 3) + (3, -2)) \in \mathcal{T}_3$. By Theorem 4.1, w is polygonal.

When $n = 2$, we will show that any word without isolated generators is polygonal. We first need a combinatorial lemma.

Lemma 4.5. *Let C be a $2m$ -gon for $m > 0$, so that each edge comes with an orientation. Label the edges of C by clean or dirty, so that clean edges and dirty edges alternate. Name each vertex v as:*

- a source, if a clean edge and dirty edge are outgoing from v ,
- a sink, if a clean edge and dirty edge are incoming to v ,
- a filter, if a dirty edge is incoming to and a clean edge is outgoing from v , and
- a pollutant, if a clean edge is incoming to and a dirty edge is outgoing from v .

Then the number of sources equals that of sinks. Also, the number of filters equals that of pollutants.

Proof. Label the vertices of C as $v_1, \dots, v_{2m}$ so that the edge (v_{2i-1}, v_{2i}) is clean for each i . Give C the orientation following $v_1, v_2, \dots, v_{2m}$. Let $h_i = 1$ if the orientation of the edge (v_i, v_{i+1}) coincides with that of C , and $h_i = -1$ otherwise. For $i = 1, \dots, 2m$, define

$$\begin{aligned}\tau(i) &= \left(\frac{1 - (-1)^i}{2}, \frac{1 + (-1)^i}{2} \right) \in \mathbb{Z}^2 \\ \sigma_i &= h_i \tau(i) - h_{i-1} \tau(i-1)\end{aligned}$$

One can check

$$\sigma_i = \begin{cases} (1, 1), & \text{if } v_i \text{ is a source;} \\ (-1, -1), & \text{if } v_i \text{ is a sink;} \\ (1, -1), & \text{if } v_i \text{ is a filter;} \\ (-1, 1), & \text{if } v_i \text{ is a pollutant.} \end{cases}$$

For example, if i is odd and v_i is a source then $\tau(i) = (1, 0), \tau(i-1) = (0, 1)$ and $h_i = 1 = -h_{i-1}$. So, $\sigma_i = (1, 0) + (0, 1) = (1, 1)$; see Figure 4. Note that $\sum_i \sigma_i = \sum_i h_i \tau(i) - \sum_i h_{i-1} \tau(i-1) = \sum_i h_i \tau(i) + \sum_i (-h_i \tau(i)) = (0, 0)$. If we let x, y, z, w be the numbers of sources, sinks, filters and pollutants, respectively, then $\sum_i \sigma_i = (x - y + z - w, x - y - z + w) = (0, 0)$. This implies that $x = y$ and $z = w$. $\square$

Theorem 4.6. *If $w \in F_2$ is a word without isolated generators, then $\rho(w) \in \mathcal{T}_2$. In particular, w is polygonal.*

Proof. We let $F_2 = \langle a, b \rangle$. Let $w = \prod_{i=1}^l a^{p_i} b^{q_i}$ for $l > 1, |p_i| > 1$ and $|q_i| > 1$. Pick two identical polygonal disks P and P' such that ∂P and $\partial P'$ read w . We consider the same partial side-pairing $\sim$ that was considered in the proof of Theorem 4.1 and put $S = P \amalg P' / \sim$. Define C to be an oriented $2l$ -gon, whose vertices are named as $v_1, \dots, v_{2l}$. Label the edges (v_{2i-1}, v_{2i}) as *clean* and the other edges as *dirty*. Also, give (v_i, v_{i+1}) the orientation coinciding that of C if $p_i > 0$, and the opposite orientation otherwise. Using the terms in Lemma 4.5, each

source v_i of C corresponds to a boundary component ∂_i of S such that there exists an outgoing a -edge and an outgoing b -edge at the two vertices of ∂_i . Following the notation in the proof of Theorem 4.1 for $a_1 = a$ and $a_2 = b$, one can write $\rho(\partial_i) = (-1, 2) = (-2, 1) \in \mathcal{A}_2$ (see Figure 4). Similarly, a sink corresponds to a boundary component ∂_i such that $\rho(\partial_i) = (1, -2) = (2, -1)$, a filter to $\rho(\partial_i) = (2, 1) = (-1, -2)$, and a pollutant to $\rho(\partial_i) = (1, 2) = (-2, -1)$. By Lemma 4.5, $\rho(w) = \sum_{i=1}^s ((1, -2) + (-2, 1)) + \sum_{i=1}^f ((1, 2) + (2, 1)) \in \mathcal{T}_2$ where s and f are the numbers of sources and filters, respectively. By Theorem 4.1, w is polygonal. $\square$




FIGURE 4. Proof of Theorem 4.6. Single and double arrows denote a -edges and b -edges, respectively.

The following theorem addresses certain cases when all the b -letters are isolated, while none of the a -letters are isolated.

Theorem 4.7. *Let $w = \prod_{i=1}^l a^{p_i} b^{q_i}$ such that $l > 1$, $|p_i| > 1$, $|q_i| = 1$, and $\sum_i (\text{sign}(p_i q_i) + \text{sign}(p_{i+1} q_i)) = 0$. Then w is polygonal.*

Proof. As in the proof of Theorem 4.6, take two identical polygonal disks P and P' reading w , rotate P' by $2\pi/|w|$, and identify the overlapping edges of P and P' whenever the label and the orientation coincide. Denote by S the bounded surface obtained by such a partial side-pairing $\sim$. Let ∂_i be the boundary component of S containing the image of the interval b^{q_i} from ∂P . Put

$$r_i = (\text{sign}(p_i), \text{sign}(q_i), \text{sign}(p_{i+1}))$$

for each $i = 1, \dots, l$. There are four types of boundary components of S as shown in Figure 5 (compare this to Figure 4). This time, we call ∂_i as

- a source if $r_i = (-1, 1, 1)$ or $(-1, -1, 1)$,
- a sink if $r_i = (1, 1, -1)$ or $(1, -1, -1)$,
- a filter if $r_i = (1, 1, 1)$ or $(-1, -1, -1)$,
- a pollutant if $r_i = (-1, 1, -1)$ or $(1, -1, 1)$.

Let x, y, z and w denote the numbers of sources, sinks, filters and pollutants, respectively. Then,

$$\begin{aligned} x &= |\{i \mid p_i p_{i+1} < 0 \text{ and } p_i < 0\}| = \sum_i \left(\frac{1 - \text{sign}(p_i p_{i+1})}{2} \right) \left(\frac{1 - \text{sign}(p_i)}{2} \right). \\ y &= |\{i \mid p_i p_{i+1} < 0 \text{ and } p_i > 0\}| = \sum_i \left(\frac{1 - \text{sign}(p_i p_{i+1})}{2} \right) \left(\frac{1 + \text{sign}(p_i)}{2} \right). \\ z &= |\{i \mid p_i p_{i+1} > 0 \text{ and } p_i q_i > 0\}| = \sum_i \left(\frac{1 + \text{sign}(p_i p_{i+1})}{2} \right) \left(\frac{1 + \text{sign}(p_i q_i)}{2} \right). \\ w &= |\{i \mid p_i p_{i+1} > 0 \text{ and } p_i q_i < 0\}| = \sum_i \left(\frac{1 + \text{sign}(p_i p_{i+1})}{2} \right) \left(\frac{1 - \text{sign}(p_i q_i)}{2} \right). \end{aligned}$$

Hence,

$$4x - 4y = \sum_i (-2\text{sign}(p_i) + 2\text{sign}(p_i^2 p_{i+1})) = 2 \sum_i (-\text{sign}(p_i) + \text{sign}(p_{i+1})) = 0$$

and

$$4z - 4w = \sum_i (2\text{sign}(p_i q_i) + 2\text{sign}(p_i^2 p_{i+1} q_i)) = 2 \sum_i (\text{sign}(p_i q_i) + \text{sign}(p_{i+1} q_i)) = 0.$$

One can identify a source with a sink (a filter with a pollutant, respectively) maintaining that the surface after the identification is still w -polygonal; each identification is described in Figure 5. Hence, there exists an extension $\sim'$ of $\sim$ such that $P \coprod P' / \sim'$ is a closed w -polygonal surface. $\square$




FIGURE 5. Proof of Theorem 4.7. If the source in (a) and the sink in (b) are glued by identifying the edges AB, BC, CD and DA with $A'B', B'C', C'D'$ and $D'A'$, respectively, the resulting surface is still w -polygonal. Similarly, the filter in (c) and the pollutant in (d) can be glued by identifying EF, FG, GH and HE with $E'F', F'G', G'H'$ and $H'E'$ respectively, without generating any vertex with two incoming or two outgoing edges of the same label.

Corollary 4.8. $\prod_{i=1}^l a^{p_{2i-1}}(a^{p_{2i}})^b$ is polygonal where $l > 0$ and $|p_i| > 1$ for each i .

Proof. Write $w = \prod_i a^{p_{2i-1}} b^{q_{2i-1}} a^{p_{2i}} b^{q_{2i}}$ where $q_{2i-1} = -1$ and $q_{2i} = 1$. Then, $\sum_i (\text{sign}(p_i q_i) + \text{sign}(p_{i+1} q_i)) = \sum_i \text{sign}(p_i)(\text{sign}(q_i) + \text{sign}(q_{i-1})) = 0$. By Theorem 4.7, w is polygonal. $\square$




5. SIMPLE HEIGHT-1 WORDS

A word $w \in F_2 = \langle a, b \rangle$ is a *simple height-one word* if for some $l > 0$, $w = \prod_{i=1}^l a^{p_i}(a^{q_i})^b$ where p_i 's are of the same sign and q_i 's are of the same sign. In particular, if $p_i, q_i > 0$, then w is called a *positive height-one word*. Note that the isomorphism $F_2 \cong \langle a, a^b \rangle$ defined by $a \mapsto a$ and $b \mapsto a^b$ maps positive words to positive height-one words. In this section, we prove polygonality for a class of simple height-one words, including the Baumslag–Solitar relator $a^p(a^q)^b$ for $pq \neq 0$. In some sense that will be made clear in Remark 5.9, *most* simple height-one words turn out to be polygonal. We will mainly focus on positive height-one words as the other cases can be dealt with in almost identical ways.

Definition 5.1. Let $w = \prod_{i=1}^l a^{p_i}(a^{q_i})^b$ be a positive height-one word. Define $p_{l+1}, p_{l+2}, \dots$ and $q_{l+1}, q_{l+2}, \dots$ by requiring $p_{i+l} = p_i$ and $q_{i+l} = q_i$ for $i > 0$. Fix $m > 0$ and consider polygonal disks $P_1, \dots, P_m$ whose boundary edges are oriented and labeled by a or b so that ∂P_i reads $w^{k_i} = \prod_{j=1}^{k_i l} a^{p_j}(a^{q_j})^b$ for some $k_i > 0$. On ∂P_i , we will denote by $\alpha_j^{(i)}$ ($\beta_j^{(i)}$, respectively) the b -edge between the intervals a^{p_j} and a^{q_j} (a^{q_j} and $a^{p_{j+1}}$, respectively); see Figure 6 (a). A partial side-pairing $\sim$ on $\coprod_i P_i$ is a *consistent b -side-pairing* if $\sim$ pairs all b -edges of $\coprod_i \partial P_i$ such that each $\alpha_j^{(i)}$ is identified with some $\beta_{j'}^{(i')}$.

We let $\mathcal{B}$ denote the free abelian monoid generated by $\lambda_{(c_1, \dots, c_r)}^+$ and $\lambda_{(c_1, \dots, c_r)}^-$ where $(c_1, \dots, c_r)$ ranges over all finite ordered tuples of positive integers. Let $\sim$ be a consistent b -side-pairing on $\coprod_i P_i$ and put $S = \coprod_i P_i / \sim$. Each boundary component ∂_i of S consists of a -edges whose orientations induce the same orientation on $\partial_i \approx S^1$. We will continue to use the notation of Definition 5.1. If $\alpha_j^{(i)}$ and $\beta_{j'}^{(i')}$ are paired by $\sim$, then some boundary component $\partial_1 \subseteq \partial S$ contains two adjacent intervals a^{p_j} from ∂P_i and $a^{p_{j'+1}}$ from $\partial P_{i'}$ which are concatenated at the terminal vertex of $\alpha_j^{(i)} \sim \beta_{j'}^{(i')}$; see Figure 6. Similarly, another boundary component $\partial_2 \subseteq \partial S$ contains the intervals $a^{q_{j'}}$ from $\partial P_{i'}$ and a^{q_j} from ∂P_i which are concatenated at the initial vertex of $\alpha_j^{(i)} \sim \beta_{j'}^{(i')}$. Note that at each boundary component ∂_i of S , b -edges are all incoming or all outgoing at their intersections with ∂_i . Let $c_1, c_2, \dots, c_r$ denote the lengths of the intervals (in this order) on ∂_i , partitioned by the vertices intersecting with b -edges. If b -edges are all incoming at ∂_i , we define $\lambda(\partial_i) = \lambda_{(c_1, \dots, c_r)}^- \in \mathcal{B}$, and if b -edges are all outgoing at ∂_i , define $\lambda(\partial_i) = \lambda_{(c_1, \dots, c_r)}^+$. For $\partial S = \partial_1 \cup \partial_2 \cup \dots$, we define $\lambda(S) = \sum_i \lambda(\partial_i)$. (This is an abuse of notation, as $\lambda(S)$ depends on the choice of the expression $S = \coprod_i P_i / \sim$ and the immersion $S^{(1)} \rightarrow \text{Cay}(F_2)/F_2$.)

Example 5.2. Consider the Baumslag–Solitar relator $w = a(a^2)^b$. By the consistent b -side-pairing $\sim$ on a polygonal disk P shown in Figure 7, $S = P / \sim$ is a three-punctured sphere with boundary components ∂_1, ∂_2 and ∂_3 . One sees that $\lambda(\partial_1) = \lambda(\partial_3) = \lambda_{(2)}^+$ and $\lambda(\partial_2) = \lambda_{(1,1)}^-$. If one takes an identical copy S' of S with each boundary component $\partial'_i \subseteq \partial S'$ corresponding to $\partial_i \subseteq \partial S$, a closed surface S_0

FIGURE 6. A consistent b -side-pairing on $P_i \amalg P_{i'}$.(a) P (b) $S = P/\sim$ FIGURE 7. (a) A polygonal disk P where ∂P reads w^2 for $w = a(a^2)^b$. Dashed curves show the consistent b -side-pairing $\sim$. (b) $S = P/\sim$ is a three-punctured sphere.

can be formed by identifying ∂_1 to ∂_2 , ∂'_1 to ∂'_2 , and ∂_3 to ∂'_3 . In order to show that S_0 is w -polygonal, one has only to be careful about the identification of ∂_3 with ∂'_3 ; that is, we identify ∂_3 to ∂'_3 by a π -rotation so that the vertex on ∂_3 to which a b -edge is outgoing (the vertex D in the Figure 7 (b)) is not identified with its copy on ∂'_3 . This implies that the Baumslag–Solitar word w is polygonal.

Definition 5.3. We let $\mathcal{U}$ be the submonoid of $\mathcal{B}$ generated by the following elements.

- $\lambda_{(c_1, \dots, c_r)}^+ + \lambda_{(d_1, \dots, d_s)}^-$ where $\sum_i c_i = \sum_j d_j$.

- $\lambda_{(c_1, \dots, c_r)}^+ + \lambda_{(d_1, \dots, d_s)}^+$ and $\lambda_{(c_1, \dots, c_r)}^- + \lambda_{(d_1, \dots, d_s)}^-$, where $m = \sum_i c_i = \sum_j d_j$ and there exists c such that the sets $\{c + c_1, c + c_1 + c_2, \dots, c + c_1 + \dots + c_r = c + m\}$ and $\{d_1, d_1 + d_2, \dots, d_1 + \dots + d_s = m\}$ are disjoint modulo m .

Lemma 5.4. *Let w be a simple height-one word. Suppose $P_1, \dots, P_k$ are polygonal disks reading some powers of w , and $\sim_i$ is a consistent b -side-pairing on P_i for each i . If $\sum_i \lambda(P_i / \sim_i) \in \mathcal{U}$, then w is polygonal.*

Proof. We will only prove the theorem for positive height-one words, as the other cases are very similar. Write $w = \prod_{i=1}^l a^{p_i} (a^{q_i})^b$ for $p_i, q_i > 0$. Let $S_i = P_i / \sim_i$ for each i . One can rewrite the sum $\sum_i \lambda(S_i) = \sum_j (\lambda(\partial_j) + \lambda(\partial'_j))$, where $\{\partial_j, \partial'_j \mid j = 1, 2, \dots\}$ is the collection of the boundary components of $\coprod_i S_i$ such that each $\lambda(\partial_j) + \lambda(\partial'_j)$ is a generator of $\mathcal{U}$. In particular, the lengths (that is, the number of edges) of ∂_j and ∂'_j are the same. We will define a pairing $\sim$ of the a -edges on $\partial(\coprod_i S_i)$ so that ∂_j and ∂'_j are identified for each j . Fix $\partial = \partial_j$ and $\partial' = \partial'_j$.

First, consider the case when $\lambda(\partial) = \lambda_{(c_1, \dots, c_r)}^+$ and $\lambda(\partial') = \lambda_{(d_1, \dots, d_s)}^-$ such that $\sum_i c_i = \sum_j d_j$. In this case, we let $\sim$ be any identification of ∂ and ∂' respecting the orientations of the a -edges. This corresponds to the identification of ∂_1 to ∂_2 and ∂'_1 to ∂'_2 in Example 5.2. Recall that at each vertex of $\partial \subseteq \coprod_i \partial S_i$ (∂' , respectively), either no b -edges are intersecting, or exactly one b -edge is outgoing (incoming, respectively). Hence, at each vertex in the image of $\partial \sim \partial'$ in $\coprod_i S_i / \sim$, no two b -edges can be both incoming or both outgoing.

Next, suppose $\lambda(\partial) = \lambda_{(c_1, \dots, c_r)}^+$ and $\lambda(\partial') = \lambda_{(d_1, \dots, d_s)}^+$; put $m = \sum_i c_i = \sum_j d_j$, and choose c such that $c + \sum_{i=1}^h c_i \not\equiv \sum_{j=1}^k d_j \pmod{m}$ for all h, k . Label the vertices of ∂ by $v_0, v_1, \dots, v_{m-1}$ consecutively, such that v_s intersects with an outgoing b -edge if and only if $s = \sum_{i=1}^h c_i$ for some $h > 0$. Moreover, we choose the labeling so that the orientation of the a -edge (v_i, v_{i+1}) is from v_i to v_{i+1} . Similarly, label the vertices of ∂' by $v'_0, v'_1, \dots, v'_{m-1}$, so that v'_s intersect with an outgoing b -edge if and only if $s = \sum_{j=1}^k d_j$ for some $k > 0$, and the orientations are from v'_i to v'_{i+1} . We let $\sim$ identify ∂ to ∂' such that v_i and v'_{i+c} are identified for $i = 0, \dots, m-1$. By the assumption, the image of ∂ in $S = \coprod_i S_i / \sim$ will not have any vertex with two outgoing b -edges. There are no incoming b -edges at ∂ and ∂' . Define $\sim$ similarly in the case when $\lambda(\partial) = \lambda_{(c_1, \dots, c_r)}^-$ and $\lambda(\partial') = \lambda_{(d_1, \dots, d_s)}^-$. As a result, we get a closed w -polygonal surface $S = (\coprod_i S_i) / \sim$. Hence, w is polygonal. $\square$

Lemma 5.5. *Let $c_1, \dots, c_r$ be positive integers.*

- (1) *If each c_i is larger than 1, then $2\lambda_{(c_1, \dots, c_r)}^\pm \in \mathcal{U}$.*
- (2) *If there exists $1 \leq i_0 \leq r$ such that $c_{i_0} \geq 1 + \frac{1}{2} \sum_{i=1}^r c_i$, then $2\lambda_{(c_1, \dots, c_r)}^\pm \in \mathcal{U}$.*

Proof. Put $m = \sum_{i=1}^r c_i$.

(1) It suffices to show that $\{c_1, c_1 + c_2, \dots, c_1 + \dots + c_r = m\}$ and $\{1 + c_1, 1 + c_1 + c_2, \dots, 1 + c_1 + \dots + c_r = 1 + m\}$ are disjoint modulo m (Definition 5.3). Assume the contrary. Then for some $1 \leq h, k \leq r$, we have $\sum_{i=1}^h c_i \equiv 1 + \sum_{i=1}^k c_i \pmod{m}$. Hence, $\sum_{i=1}^h c_i - \sum_{i=1}^k c_i \equiv 1 \pmod{m}$; that is, the sum of some consecutive terms in $c_1, c_2, \dots, c_r$ is congruent to ± 1 modulo m . This is a contradiction, since $c_i > 1$ for each i .

(2) Without loss of generality, let $i_0 = 1$. We have $c_1 \geq 1 + \frac{1}{2}m$. In particular, $c_1 \geq 2 + m - c_1 \geq 2$. We claim that $A = \{\sum_{i=1}^h c_i \mid 1 \leq h \leq r\}$ and $B =$

$\{(m+1-c_1) + \sum_{i=1}^k c_i \mid 1 \leq k \leq r\}$ are disjoint modulo m . Suppose $\sum_{i=1}^h c_i \equiv m+1-c_1 + \sum_{i=1}^k c_i \pmod{m}$ for some $1 \leq h, k \leq r$. If $h \geq k$ then it follows that $c_1 + \sum_{i=k+1}^h c_i \equiv 1 \pmod{m}$, which is impossible since $2 \leq c_1 + \sum_{i=k+1}^h c_i \leq \sum_{i=1}^r c_i = m$. If $h < k$, then $c_1 \equiv 1 + \sum_{i=h+1}^k c_i \pmod{m}$, which is also impossible since $1 + \sum_{i=h+1}^k c_i \leq 1 + \sum_{i=2}^r c_i = 1 + (m - c_1) \leq 1 + (c_1 - 2) = c_1 - 1$. $\square$

In Example 5.2, $\lambda(S) = \lambda(\partial_1) + \lambda(\partial_2) + \lambda(\partial_3) = 2\lambda_{(2)}^- + \lambda_{(1,1)}^+$. So, $\lambda(S) + \lambda(S) = 4\lambda_{(2)}^- + 2\lambda_{(1,1)}^+ = 2\lambda_{(2)}^- + 2(\lambda_{(2)}^- + \lambda_{(1,1)}^+) \in \mathcal{U}$. By Lemma 5.4, we prove again that $w = a(a^2)^b$ is polygonal.

Theorem 5.6. *Let $w = \prod_{i=1}^l a^{p_i}(a^{q_i})^b$ be a simple height-one word. Put $p = \sum_{i=1}^l |p_i|$ and $q = \sum_{i=1}^l |q_i|$. Let p' and q' denote the number of elements in $\{i \mid p_i = \pm 1\}$ and $\{i \mid q_i = \pm 1\}$, respectively. If $pp' \leq q^2$ and $qq' \leq p^2$, then w is polygonal.*

Proof. We will give a proof only for the case when w is a positive height-one word as the other cases are very similar. Without loss of generality, we may assume that $pp' \geq qq'$. Let c be the least common multiple of $\{q_1, q_2, \dots, q_l\}$ and put $d = c!$. Let $P_1, P_2, \dots, P_c$ ($Q_1, Q_2, \dots, Q_c$, respectively) be polygonal disks reading w^{dp} (w^{dq} , respectively). Define $p_{l+1}, p_{l+2}, \dots$ and $q_{l+1}, q_{l+2}, \dots$ by $p_{i+l} = p_i$ and $q_{i+l} = q_i$ for each i . Especially, $w^k = \prod_{i=1}^{kl} a^{p_i}(a^{q_i})^b$ for any $k > 0$. Following the notations in Definition 5.1, we let $\alpha_j^{(i)}$ ($\beta_j^{(i)}$, respectively) denote the b -edge between intervals a^{p_j} and a^{q_j} (a^{q_j} and $a^{p_{j+1}}$, respectively) on ∂P_i . Similarly, let $\gamma_j^{(i)}$ ($\delta_j^{(i)}$, respectively) be the b -edge between a^{p_j} and a^{q_j} (a^{q_j} and $a^{p_{j+1}}$, respectively) on ∂Q_i . For convention, we let $\beta_0^{(i)} = \beta_{dpl}^{(i)}$ and $\delta_0^{(i)} = \delta_{dql}^{(i)}$. Define a consistent b -side-pairing $\sim$ on $(\coprod_i P_i) \coprod (\coprod_i Q_i)$ by

- $\beta_{j-1}^{(i)} \sim \alpha_j^{(i)}$ for each $1 \leq i \leq c$ and $1 \leq j \leq dpl$, and
- $\gamma_j^{(i)} \sim \delta_j^{(i)}$ for each $1 \leq i \leq c$ and $1 \leq j \leq dql$.

As is easily seen in the example shown in Figure 8 (a) and (b),

$$\begin{aligned} \lambda(P_i/\sim) &= \sum_{j=1}^{dpl} \lambda_{(p_j)}^- + \lambda_{(q_1, q_2, \dots, q_{dpl})}^+ \\ &= dpp' \lambda_{(1)}^- + d \sum_{\substack{1 \leq j \leq pl \\ p_j > 1}} \lambda_{(p_j)}^- + \lambda_{(q_1, q_2, \dots, q_{dpl})}^+ \\ \lambda(Q_i/\sim) &= dq q' \lambda_{(1)}^+ + d \sum_{\substack{1 \leq j \leq ql \\ q_j > 1}} \lambda_{(q_j)}^+ + \lambda_{(p_1, p_2, \dots, p_{dql})}^- \end{aligned}$$

Let $r = pp' - qq'$. First, consider the case when $r = 0$. Put $S = (\coprod_{i=1}^c P_i) \coprod (\coprod_{i=1}^c Q_i)/\sim$.

$$\begin{aligned} 2\lambda(S) &= 2cdpp'(\lambda_{(1)}^- + \lambda_{(1)}^+) + cd \sum_{\substack{1 \leq j \leq pl \\ p_j > 1}} 2\lambda_{(p_j)}^- + cd \sum_{\substack{1 \leq j \leq ql \\ q_j > 1}} 2\lambda_{(q_j)}^+ \\ &\quad + 2c \left(\lambda_{(q_1, q_2, \dots, q_{dpl})}^+ + \lambda_{(p_1, p_2, \dots, p_{dql})}^- \right) \end{aligned}$$

Since $q_1 + q_2 + \dots + q_{dpl} = dp(q_1 + q_2 + \dots + q_l) = dpq = p_1 + p_2 + \dots + p_{dql}$, the definition of $\mathcal{U}$ and Lemma 5.5 imply that $2\lambda(S) \in \mathcal{U}$. By Lemma 5.4, w is polygonal.

Now assume that $r > 0$. Since $r \leq pp' = |\{j \mid 1 \leq j \leq pl, p_j = 1\}|$, one can choose $A \subseteq \{j \mid 1 \leq j \leq pl, p_j = 1\}$ such that $|A| = r$. Find $x_1, \dots, x_l \geq 0$ satisfying that

- (i) $x_j \leq qq_j$ and $\sum_j x_j = r$,
- (ii) if $q_j = 1$, then $x_j = 0$.

Such x_j 's exist, since

$$\sum_{\substack{1 \leq j \leq l \\ q_j \neq 1}} qq_j = q \left(\sum_{1 \leq j \leq l} q_j - \sum_{\substack{1 \leq j \leq l \\ q_j = 1}} q_j \right) = q(q - q') \geq pp' - qq' = r$$

Fix a map $\sigma: A \rightarrow \{1, \dots, l\}$ such that $|\sigma^{-1}(j)| = x_j$. We will modify $\sim$ to obtain another consistent b -side-pairing $\sim'$ on $(\coprod_i P_i) \coprod (\coprod_i Q_i)$. On $\coprod_i \partial Q_i$, we require that $\sim'$ coincide with $\sim$; that is, $\gamma_j^{(i)} \sim' \delta_j^{(i)}$ for each $1 \leq i \leq c$ and $1 \leq j \leq dql$. We define $\sim'$ on $\coprod_i \partial P_i$ as follows. For $1 \leq i \leq c$ and $1 \leq j \leq dpl$,

- $\beta_{j-1}^{(i)} \sim' \alpha_j^{(i+1)}$, if $j \equiv j' \pmod{pl}$ for some $j' \in A$ and $i \not\equiv 0 \pmod{q_{\sigma(j')}}$,
- $\beta_{j-1}^{(i)} \sim' \alpha_j^{(i+1-q_{\sigma(j')})}$, if $j \equiv j' \pmod{pl}$ for some $j' \in A$ and $i \equiv 0 \pmod{q_{\sigma(j')}}$,
- $\beta_{j-1}^{(i)} \sim' \alpha_j^{(i)}$, if $j \not\equiv j' \pmod{pl}$ for any $j' \in A$.

Put $S' = (\coprod_i P_i) \coprod (\coprod_i Q_i) / \sim'$. For each $k \in A$, define $g_k \in \text{SYM}(\{1, 2, \dots, c\})$ by

$$g_k(i) = \begin{cases} i+1, & \text{if } i \not\equiv 0 \pmod{q_{\sigma(k)}} \\ i+1-q_{\sigma(k)}, & \text{otherwise.} \end{cases}$$

Then the order of g_k is $q_{\sigma(k)}$. For an arbitrary $1 \leq i_0 \leq c$, define $i_1, i_2, \dots$ by

$$i_j = \begin{cases} g_{j'}(i_{j-1}), & \text{if } j \equiv j' \pmod{pl} \text{ for some } j' \in A \\ i_{j-1}, & \text{otherwise} \end{cases}$$

Then $\beta_{j-1}^{(i_{j-1})} \sim' \alpha_j^{(i_j)}$ for $1 \leq j \leq dpl$. Since $d = cl$, $(\prod_{k \in A} g_k)^d = \text{Id} \in \text{SYM}(\{1, 2, \dots, c\})$; hence, $i_{dpl} = (\prod_{k \in A} g_k)^d(i_0) = i_0$. So, $\alpha_{dpl}^{(i_{dpl})} = \alpha_{dpl}^{(i_0)}$. Pick any $1 \leq i_0 \leq c$ and denote by ∂_{i_0} the boundary component of S' containing the interval $a^{q_{dpl}}$ from $\partial P_{i_0} = \partial P_{i_{dpl}}$. Since the interval $a^{q_{dpl}}$ from ∂P_{i_0} terminates at the initial vertex of $\beta_{dpl}^{(i_0)} = \beta_0^{(i_0)}$, and $\beta_0^{(i_0)} \sim' \alpha_1^{(i_1)}$, the initial vertex of a^{q_1} from ∂P_{i_1} is concatenated at the terminal vertex of $a^{q_{dpl}}$ from ∂P_{i_0} , in ∂_{i_0} . In this way, one sees that ∂_{i_0} contains the interval a_{q_j} from ∂P_{i_j} for each $1 \leq j \leq dpl$; for example, see Figure 8 (c). It follows that $\lambda(\partial_{i_0}) = \lambda_{(q_1, q_2, \dots, q_{dpl})}^+$. On the other hand, if $1 \leq j \leq dpl$ and $j \equiv j' \pmod{pl}$ for some $j' \in A$, then the terminal vertex of the a -edge corresponding to a^{p_j} on ∂P_i is concatenated with the initial vertex of a^{p_j} on $\partial P_{g_j^{-1}(i)}$; in particular, the a -edges corresponding to a^{p_j} on $\partial P_1, \partial P_2, \dots, \partial P_c$ form boundary components $\partial'_1, \partial'_2, \dots, \partial'_{c/q_{\sigma(j')}}$ in S' where for each $i = 1, 2, \dots, c/q_{\sigma(j')}$,

$$\lambda(\partial'_i) = \underbrace{\lambda_{(1, \dots, 1)}^-}_{q_{\sigma(j')}}.$$

Recall that

$$\lambda(\coprod_i P_i / \sim) = cdpp' \lambda_{(1)}^- + cd \sum_{\substack{1 \leq j \leq pl \\ p_j > 1}} \lambda_{(p_j)}^- + c \lambda_{(q_1, q_2, \dots, q_{dpl})}^+$$

We have

$$\begin{aligned}
\lambda(\coprod_i P_i/\sim') &= cd \sum_{\substack{1 \leq j \leq pl \\ p_j=1, j \notin A}} \lambda_{(1)}^- + d \sum_{j \in A} \left(\frac{c}{q_{\sigma(j)}} \right) \lambda_{(\underbrace{1, \dots, 1}_{q_{\sigma(j)}})}^- \\
&\quad + cd \sum_{\substack{1 \leq j \leq pl \\ p_j > 1}} \lambda_{(p_j)}^- + c \lambda_{(q_1, q_2, \dots, q_{dpl})}^+ \\
&= cd(pp' - r) \lambda_{(1)}^- + \sum_{\substack{1 \leq j \leq l \\ q_j > 1}} dx_j \left(\frac{c}{q_j} \right) \lambda_{(\underbrace{1, \dots, 1}_{q_j})}^- \\
&\quad + cdp \sum_{\substack{1 \leq j \leq l \\ p_j > 1}} \lambda_{(p_j)}^- + c \lambda_{(q_1, q_2, \dots, q_{dpl})}^+. \\
\lambda(\coprod_i Q_i/\sim') &= \lambda(\coprod_i Q_i/\sim) = cdq q' \lambda_{(1)}^+ + cdq \sum_{\substack{1 \leq j \leq l \\ q_j > 1}} \lambda_{(q_j)}^+ + c \lambda_{(p_1, p_2, \dots, p_{dql})}^+
\end{aligned}$$

Note that $pp' - r = qq'$. From $\lambda(S') = \lambda(\coprod_i P_i/\sim') + \lambda(\coprod_i Q_i/\sim')$, we have:

$$\begin{aligned}
2\lambda(S') &= 2cdq q' (\lambda_{(1)}^- + \lambda_{(1)}^+) + \sum_{\substack{1 \leq j \leq l \\ q_j > 1}} 2dx_j \left(\frac{c}{q_j} \right) (\lambda_{(\underbrace{1, \dots, 1}_{q_j})}^- + \lambda_{(q_j)}^+) \\
&\quad + \sum_{\substack{1 \leq i \leq l \\ q_j > 1}} \left(cdq - dx_j \left(\frac{c}{q_j} \right) \right) 2\lambda_{(q_j)}^+ \\
&\quad + cdp \sum_{\substack{1 \leq j \leq l \\ p_j > 1}} 2\lambda_{(p_j)}^- + c \left(2\lambda_{(q_1, q_2, \dots, q_{dpl})}^+ + 2\lambda_{(p_1, p_2, \dots, p_{dql})}^- \right).
\end{aligned}$$

Note that $cdq - dx_i(c/q_i) \geq 0$ by the choice of x_i . Hence, $2\lambda(S') \in \mathcal{U}$. In particular, w is polygonal. $\square$

- Example 5.7.** (1) Let us illustrate the proof of Theorem 5.6 for the Baumslag-Solitar relator $w = a(a^2)^b$. Note $p = 1 = p', q = 2, q' = 0$ and $c = 2 = d$. As in Figure 8 (a) and (b), choose P_1, P_2, Q_1, Q_2 such that ∂P_i reads $w^{cp} = w^2$ and ∂Q_i reads $w^{cq} = w^4$. On P_i and Q_i , a consistent b -side-pairing $\sim$ is shown by dashed lines. Then, $\lambda(P_1/\sim) = \lambda(P_2/\sim) = 2\lambda_{(1)}^- + \lambda_{(2,2)}^+$ and $\lambda(Q_1/\sim) = \lambda(Q_2/\sim) = \lambda_{(1,1,1,1)}^- + 4\lambda_{(2)}^+$. Let a new partial side-pairing $\sim'$ coincide with $\sim$ on $\coprod \partial Q_i$ and be modified on $\coprod \partial P_i$ as shown in Figure 8 (c). We have $\lambda(P_1 \coprod P_2/\sim') = 2\lambda_{(1,1)}^- + 2\lambda_{(2,2)}^+$. Define $S' = (\coprod_i P_i) \coprod (\coprod_i Q_i)/\sim'$. $2\lambda(S') = 4\lambda_{(1,1)}^- + 4\lambda_{(2,2)}^+ + 4\lambda_{(1,1,1,1)}^- + 16\lambda_{(2)}^+ = 4(\lambda_{(1,1)}^- + \lambda_{(2)}^+) + 4(\lambda_{(2,2)}^+ + \lambda_{(1,1,1,1)}^-) + 6(2\lambda_{(2)}^+) \in \mathcal{U}$.
- (2) More generally, any Baumslag-Solitar relator $w = a^p(a^q)^b$ is a simple height-one word for $pq \neq 0$. As pp' and qq' are either 0 or 1, $pp' \leq 1 \leq q^2$ and $qq' \leq 1 \leq p^2$. By Theorem 5.6, w is polygonal.

Corollary 5.8. Let $w = \prod_{i=1}^l a^{p_i}(a^{q_i})^b$ be a simple height-one word. Put $p = \sum_{i=1}^l |p_i|$ and $q = \sum_{i=1}^l |q_i|$. If there exists $r > 1$ such that $\min(p/l, q/l) \geq r$ and $1/r \leq p/q \leq r$, then w is polygonal.




FIGURE 8. Example 5.7 (1). The orientations and the labels of the edges of ∂P_i and ∂Q_i . In (a) and (c), the vertices labeled by the same alphabets are actually the same in P_i .

Proof. Note that $p' = \sum_{|p_i|=1} 1 \leq l$, and similarly, $q' = \sum_{|q_i|=1} 1 \leq l$. $pp' \leq pl \leq rq \cdot (q/r) = q^2$, and by symmetry, $qq' \leq p^2$. By Theorem 5.6, w is polygonal. $\square$

Remark 5.9. Let N be a sufficiently large integer. Consider the probability space X_N of the positive words of length N in $F_2 = \langle a, b \rangle$, where a and b are equally likely to appear. Set f to be the injective endomorphism on F_2 defined by $f(a) = a$ and $f(b) = a^b$. Let $Y_N = f(X_N)$ have the push-forward probability from X_N by f . Write an arbitrary word $w \in Y_N$ as $w = f(\prod_{i=1}^m a^{c_i} b^{d_i}) = \prod_{i=1}^m a^{c_i} (a^{d_i})^b$, where $m > 0$, $c_i, d_i \geq 0$ and moreover, $c_i, d_{m+1-i} \neq 0$ unless $i = 1$. Let s denote the number of *runs* in w ; this means, s is the number of non-trivial terms in $(a^{c_1}, b^{d_1}, \dots, a^{c_m}, b^{d_m})$. If $w = a^N$ or $w = (a^N)^b$, we put $(p_1, q_1) = (N, 0)$ or $(p_1, q_1) = (0, N)$ respectively, and also let $w' = w$ and $l = \frac{1}{2}$; otherwise, find a cyclic conjugation $w' \sim w$ such that $w' = \prod_{i=1}^l a^{p_i} (a^{q_i})^b$ where $p_i, q_i > 0$ for all i . Define $p = \sum_i p_i, q = \sum_i q_i, p' = \sum_{|p_i|=1} 1$ and $q' = \sum_{|q_i|=1} 1$. Note that p, q, p', q', l and s are well-defined by w , and that p and q have the binomial distribution $B(N, 1/2)$. It is obvious that $p', q' \leq l \leq \frac{1}{2}s$ and so,

$$P(q^2 \leq pp') \leq P(q^2 \leq \frac{1}{2}ps).$$

Let us set $\epsilon = \frac{1}{6}$ and $\alpha = ((1 - \epsilon)/(1 + \epsilon))^2 = 25/49 > 1/2$. Denote by A the event that $\frac{1-\epsilon}{2}N \leq q \leq \frac{1+\epsilon}{2}N$. If A occurs, then $2q^2/p = \frac{q}{N-q} \cdot 2q \geq \frac{1-\epsilon}{1+\epsilon} \cdot (1 - \epsilon)N$. So,

$$\begin{aligned} P(q^2 \leq \frac{1}{2}ps) &= P((2q^2/p \leq s) \text{ and } A) + P((2q^2/p \leq s) \text{ and } A^c) \\ &\leq P\left(\left(\frac{(1-\epsilon)^2}{1+\epsilon}N \leq s\right) \text{ and } A\right) + P(A^c) \\ &\leq P\left(s \geq \frac{(1-\epsilon)^2}{1+\epsilon}N\right) + P(A^c) \leq P(s-1 \geq \alpha(N-1)) + P(A^c). \end{aligned}$$

The last inequality follows from the fact that

$$\frac{(1-\epsilon)^2}{1+\epsilon}N - 1 \geq \left(\frac{1-\epsilon}{1+\epsilon}\right)^2 (N-1) = \alpha(N-1)$$

for sufficiently large N . Since $s-1$ has the binomial distribution $B(N-1, 1/2)$ [7], the expectation and the standard deviation of $s-1$ are $(N-1)/2$ and $\sqrt{N-1}/2$, respectively. By Chebyshev's inequality,

$$\begin{aligned} P(s-1 \geq \alpha(N-1)) &= P\left(s-1 - \frac{N-1}{2} \geq (\alpha - \frac{1}{2})(N-1)\right) \\ &= P\left(\frac{s-1 - (N-1)/2}{\sqrt{N-1}/2} \geq 2(\alpha - \frac{1}{2})\sqrt{N-1}\right) \leq \frac{1}{(2\alpha - 1)^2(N-1)} \end{aligned}$$

and

$$\begin{aligned} P(A^c) &= P(|q - N/2| > \epsilon N/2) \\ &= P\left(\frac{|q - N/2|}{\sqrt{N}/2} > \epsilon\sqrt{N}\right) \leq \frac{1}{\epsilon^2 N} \leq \frac{1}{\epsilon^2(N-1)}. \end{aligned}$$

Therefore, $P(q^2 \leq pp') \leq C/(N-1)$ for some constant $C > 0$. In particular, $P(q^2 \leq pp')$ converges to 0 as N approaches infinity. By symmetry, $P(p^2 \leq qq') \rightarrow 0$ as $N \rightarrow \infty$. From Theorem 5.6, it follows that the probability for a word in Y_N to be polygonal converges to 1 as $N \rightarrow \infty$. The same argument applies to the cases when $c_i < 0$ or $d_i < 0$. In this sense, one concludes that simple height-one words are *almost surely* polygonal.

REFERENCES

- [1] M Bestvina and M Feighn. A combination theorem for negatively curved groups. *J. Differential Geom.*, 35(1):85–101, 1992.
- [2] Martin R. Bridson and André Haefliger. *Metric spaces of non-positive curvature*, volume 319 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1999.
- [3] D Calegari. Surface subgroups from homology. *Geometry & Topology*, 12(4):1995–2007, 2008.
- [4] C McA Gordon and H Wilton. On surface subgroups of doubles of free groups. *J. Lond. Math. Soc. (2)*, 82(1):17–31, 2010.
- [5] Frédéric Haglund. Finite index subgroups of graph products. *Geom. Dedicata*, 135:167–209, 2008.
- [6] Jon McCammond. Personal communication, 2010.
- [7] A. M Mood. The distribution theory of runs. *Ann. Math. Statistics*, 11:367–392, 1940.
- [8] P Scott. Subgroups of surface groups are almost geometric. *J. London Math. Soc. (2)*, 17(3):555–565, 1978.
- [9] P Scott and T Wall. Topological methods in group theory. *Homological group theory (Proc. Sympos., Durham, 1977)*, 137–203, London Math. Soc. Lecture Note Ser., 36, Cambridge Univ. Press, Cambridge-New York, 1979.

- [10] J R Stallings. Topology of finite graphs. *Invent. Math.*, 71(3):551–565, 1983.
- [11] Henry Wilton. Hall’s theorem for limit groups. *Geom. Funct. Anal.*, 18(1):271–303, 2008.
- [12] D T Wise. Subgroup separability of graphs of free groups with cyclic edge groups. *Q. J. Math.*, 51(1):107–129, 2000.

DEPARTMENT OF MATHEMATICAL SCIENCES, KAIST, 335 GWAHANGNO, YUSEONG-GU, DAEJEON
305-701, REPUBLIC OF KOREA

E-mail address: `shkim@kaist.edu`

MATHEMATICS 253-37, CALTECH, PASADENA, CA 91125, USA

E-mail address: `wilton@caltech.edu`